

# A Categorification of Biquandle Brackets

Adu Vengal Vilas Winstein

Mathematics Department, the Ohio State University, Advised by Prof. Sergei Chmutov


## Biquandles

A *biquandle* is a set  $X$  with two binary operations  $\triangleright, \bar{\triangleright}$  such that for every  $x, y, z \in X$ ,

- $x \triangleright x = x \bar{\triangleright} x$
- The maps  $\alpha_y(x) = x \bar{\triangleright} y$ ,  $\beta_y(x) = x \triangleright y$ , and  $S(x, y) = (y \bar{\triangleright} x, x \triangleright y)$  are invertible.
- The following exchange laws are satisfied:

$$\begin{aligned} (x \triangleright y) \triangleright (z \triangleright y) &= (x \triangleright z) \triangleright (y \bar{\triangleright} z) \\ (x \triangleright y) \bar{\triangleright} (z \triangleright y) &= (x \bar{\triangleright} z) \triangleright (y \bar{\triangleright} z) \\ (x \bar{\triangleright} y) \bar{\triangleright} (z \bar{\triangleright} y) &= (x \bar{\triangleright} z) \bar{\triangleright} (y \triangleright z). \end{aligned}$$

The biquandle axioms mirror the three Reidemeister moves for link diagrams. For example, if the strands of a link are colored by elements of a biquandle so that the following relations hold at each crossing:



then the biquandle axioms are exactly what is required to allow such a coloring to be invariant under the Reidemeister moves.


Examples of biquandles include

- The trivial biquandle:*  $X = \{x\}$  is a singleton, with operations  $x \bar{\triangleright} x = x \triangleright x = x$ .
- Constant action biquandles:*  $X$  is any set,  $\sigma : X \rightarrow X$  is any bijection. The operations are  $x \triangleright y = x \bar{\triangleright} y = \sigma(x)$  for all  $x, y \in X$ .
- Alexander biquandles:*  $X$  is any module over  $\mathbb{Z}[t^{\pm 1}, r^{\pm 1}]$ . Then  $x \triangleright y = tx + (r - t)y$  and  $x \bar{\triangleright} y = ry$  define a biquandle.

Given any biquandle  $X$ , the number of  $X$ -colorings of a link diagram is an invariant and is called the *biquandle counting invariant*. If  $L$  is a link, the  $X$ -counting invariant of  $L$  is  $\Phi_X^{\mathbb{Z}}(L)$ .


## Biquandle Brackets

Let  $X$  be a biquandle and  $R$  a commutative unital ring. We would like to choose elements  $A_{x,y}, B_{x,y} \in R^{\times}$  (for each  $x, y \in X$ ),  $w \in R^{\times}$ , and  $\delta \in R$  such that the element of  $R$  determined by the following skein relations:



with  $\delta$  the value of a circle and  $w$  the value of a positive kink, is an invariant of  $X$ -colored links. Such a collection of elements of  $R$  is called a *biquandle bracket* and must satisfy three axioms found in [1].

It turns out that that  $w$  and  $\delta$  are determined by  $A$  and  $B$ . So a biquandle bracket is denoted  $\beta = (A, B)$  and its value on a coloring  $f$  of a link is denoted  $\beta(f)$ . Here is an example computation:



$$\beta(f) = w^{-3} \begin{pmatrix} +\delta B_{x,y} A_{y,z} A_{z,x} + \delta^2 B_{x,y} B_{y,z} A_{z,x} \\ +\delta A_{x,y} A_{y,z} A_{z,x} + \delta A_{x,y} B_{y,z} A_{z,x} + \delta^2 B_{x,y} A_{y,z} B_{z,x} + \delta^3 B_{x,y} B_{y,z} B_{z,x} \\ +\delta A_{x,y} A_{y,z} B_{z,x} + \delta^2 A_{x,y} B_{y,z} B_{z,x} \end{pmatrix}$$

The following multiset-valued function on links is a link invariant which enhances  $\Phi_X^{\mathbb{Z}}(L)$ :

$$\Phi_X^{\beta}(L) = \{\beta(f) : f \text{ is an } X\text{-coloring of } L\}.$$

When  $X = \{x\}$  is the trivial biquandle, we can let  $R = \mathbb{Z}[q^{\pm 1}]$ , and let  $A_{x,x} = q$  and  $B_{x,x} = q^{-1}$  to obtain the Jones polynomial as the single element of  $\Phi_X^{\beta}(L)$ . Additionally, using a computer, one can find many other biquandles and many many other biquandle brackets.

## Our Construction


Let  $\beta = (A, B)$  be an  $X$ -bracket with values in  $R$ .

- Let  $q_{x,y} = -\frac{B_{x,y}}{A_{x,y}}$  for all  $x, y \in X$ .
- Let  $x_0 \in X$ , and let  $q = q_{x_0, x_0}$ .
- Let  $G$  be the group  $\langle qq_{x,y}^{-1} : x, y \in X \rangle \leq R^{\times}$ .
- Let  $S$  be the  $R^{\times}$ -graded group algebra  $\mathbb{Z}[G]$ , with the  $R^{\times}$ -grading given by  $\deg(g) = g$  for all  $g \in G$ .
- Let  $M$  be the  $R^{\times}$  graded  $S$ -module  $S[t]/(t^2)$  with the additional grading given by  $\deg(1) = q$  and  $\deg(t) = q^{-1}$ .  $M$  is a Frobenius algebra with the following multiplication and comultiplication:

$$\begin{aligned} m : M \otimes M &\rightarrow M \\ m : 1 \otimes 1 &\mapsto 1, & 1 \otimes t &\mapsto t, \\ t \otimes 1 &\mapsto t, & t \otimes t &\mapsto 0 \\ \Delta : M &\rightarrow M \otimes M \\ \Delta : 1 &\mapsto 1 \otimes t + t \otimes 1, & t &\mapsto t \otimes t \end{aligned}$$


- Let  $L$  be a link and let  $f$  be an  $X$ -coloring of  $L$ . Perform splittings as in the figure to the left.

- Replace circles with copies of  $M$  and tensor adjacent copies together. Add maps between the modules as follows:



This gives a cube with anti-commutative faces.

- Sum the modules and maps along the columns:




to obtain  $\tilde{C}_{\beta}(f)$ , an  $R^{\times}$  graded cochain complex.

## Our Construction, Continued

- Let  $C_{\beta}(f)$  be the shifted cochain complex:
 
$$C_{\beta}(f) = \tilde{C}_{\beta}(f)[n_{-}]\{(-1)^{n-w}w^{-n+w}n_{-}\},$$
 where  $n_{\pm}$  is the number of positive or negative crossings in the link.
- Finally, take the cohomology of  $C_{\beta}(f)$  to obtain a complex  $\mathcal{H}_{\beta}(f)$ . The multiset
 
$$Bh_{\beta}(L) = \{\mathcal{H}_{\beta}(f) : f \text{ is an } X\text{-coloring of } L\}$$
 is an invariant of links.

## Results

The biquandle bracket generalizes the Jones polynomial, which Khovanov's homology construction [2] categorifies. Our goal was to generalize Khovanov's construction to biquandle brackets and obtain  $\star$ :



The Euler characteristic of  $Bh_{\beta}(L)$  is actually  $\text{rdim}(S) \cdot \Phi_X^{\beta}(L)$ , where  $\text{rdim}(S)$  is the  $R^{\times}$ -graded dimension of  $S$  which cannot always be factored out to yield  $\Phi_X^{\beta}(L)$ . In fact,  $Bh_{\beta}(L)$  is actually isomorphic to a quotient of  $Kh(L)$  shifted by

$$\text{rdim}(S) \left( \prod_{\tau^+} A_{x,y} A_{x_0, x_0}^{-1} \right) \left( \prod_{\tau^-} B_{x,y}^{-1} B_{x_0, x_0} \right).$$

This shift is the value of a *biquandle 2-cocycle invariant* which takes values in  $R^{\times}/G$ . Further work will explore how this new invariant compares to the related biquandle bracket.

## References

- Sam Nelson, Michael E Orrison, and Veronica Rivera. Quantum enhancements and biquandle brackets. *Journal of Knot Theory and Its Ramifications*, 26(05):1750034, 2017.
- Mikhail Khovanov. A categorification of the jones polynomial. *Duke Mathematical Journal*, 101(3):359–426, 2000.