



# The geometry of Ford Circles



Well  
approximable  
by  $\frac{p}{q}$ .



$$\text{Want } \delta(x, x') = \frac{k}{q^2} + \frac{k'}{q'^2} = \\ \text{in negeney: } \parallel$$

$$\sqrt{\left(\frac{p}{q} - \frac{p'}{q'}\right)^2 + \left(\frac{k}{q^2} - \frac{k'}{q'^2}\right)^2}$$

$$\Rightarrow (pq - pq')^2 = 4k^2$$

VI

|

if  $k = \frac{1}{2}$ , circles are tangent iff  $\begin{vmatrix} p & q \\ p' & q' \end{vmatrix} = \pm 1$ .

and are external o.w.

So circles tessellate.

Def:  $C_{p/q}$  is ford circle on  $\frac{p}{q}$ .

$\sim \sim \sim C_{p/q}$  tangent iff  $\begin{vmatrix} p & q \\ p' & q' \end{vmatrix} = \pm 1$ , o.w. no,

Proof:  $C_{\frac{p}{q}}$  and  $C_{\frac{p'}{q'}}$  are tangent iff  $\begin{vmatrix} p & p' \\ q & q' \end{vmatrix} = \pm 1$ , o.w. are wholly external.


Def:  $\frac{p}{q}, \frac{p'}{q'}$  are adjacent  $(\frac{p}{q} | \frac{p'}{q'})$  if  $C_{\frac{p}{q}}$  and  $C_{\frac{p'}{q'}}$  are tangent.

Theorem: If  $\frac{p}{q} \mid \frac{P_0}{Q_0}$  then all the other fractions adj to  $\frac{p}{q}$  are

$$\frac{P_n}{Q_n} = \frac{P_0 + nP}{Q_0 + nq}$$

Proof  $\frac{P_n}{Q_n} \mid \frac{p}{q}$  since  $P_n q - Q_n p = P_0 q + nqp - Q_0 p - nqr = \pm 1$

$\frac{P_n}{Q_n} \mid \frac{P_{n+1}}{Q_{n+1}}$  since  $(P_0 + np)(Q_0 + (n+1)q) - (Q_0 + nq)(P_0 + (n+1)p) = \pm 1$



$$\frac{P_n}{Q_n} - \frac{p}{q} \xrightarrow[n \rightarrow \infty]{\text{vs}} 0 \text{ from right \& left} \quad \xrightarrow[n \rightarrow -\infty]{} 0$$

no more room to fit more circles

If  $\frac{p}{q} \mid \frac{p'}{q'}$  then the mediant  $\frac{p''}{q''}$  is the unique

$$\frac{p}{q} \mid \frac{p''}{q''} \mid \frac{p'}{q'} \quad \frac{p''}{q''} = \frac{p + p'}{q + q'}$$

$\frac{p_1}{q_1} = \frac{0}{1}$ ,  $\frac{P_0}{Q_0} = \frac{1}{0}$ . All adjacencies to  $\frac{p_1}{q_1}$  are  $\frac{1}{n}$



adjacent


given  $[0; a_1, \dots]$ , then  $\frac{p_2}{q_2} = \frac{1}{a_1}$ ,  $\frac{p_3}{q_3} = \frac{p_1 + n p_2}{q_1 + n q_2}$ .

Let  $n = a_2 \Rightarrow \frac{p_3}{q_3} = \frac{a_2 p_2 + p_1}{a_2 q_2 + q_1}, \dots$



depending on parity of  $n$ .  
count cw or ccw, go around  
an times

Theorem If  $L$  is a path above the real line, and  $L$  intersects  $C_{\frac{p}{q}}$  and  $C_{\frac{p'}{q'}}$ , then  $\frac{p}{q} \mid \frac{p'}{q'}$ . (in succession)



Def A farey sequence  $F_n$  is the collection  $\left\{ \frac{p}{q} \in \mathbb{Q}, q \leq n \right\}$

take  $L$  to be a horizontal line  $y = \alpha$ .

the circles intersected are a farey sequence.

Cor. Consec. terms in  $F_n$  are adjacent.

Cor anything added from  $F_n$  to  $F_{n+1}$  is a mediant  
of 2 consec. elements of  $F_n$ .

if  $L$  is  $y = mx$  and  $L$  intersects  $r_1, \dots, r_n$

then  $\{r_1, \dots, r_n\} = \left\{ \frac{p}{q} \in \mathbb{Q} : pq \leq n \right\}$ , for some  $n(m)$ .

||  
 $F_n$  (generalized) Farey

Theorem: if  $\alpha \notin \mathbb{Q}$  then  $\left| \alpha - \frac{p}{q} \right| < \frac{1}{2q^2}$  has infinitely many solutions,

take  $L$  to be  $x = \alpha$ .


Thm: if  $L$  crosses a mesh triangle, then one of the rationals whose circle bounds the triangle satisfies

$$\left| \alpha - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}$$


Proof: Let  $\frac{P}{Q}, \frac{p}{q}, \frac{p_i}{q_i}$  bound a mesh triangle that

$L$  intersects. Say  $Q \leq q$ ,  $p_i = p + P$ ,  $q_i = q + Q$ ,

and  $\frac{P}{Q} > \frac{p}{q}$  (can get here by reflecting), so that  $Pq - Qp = 1$



Solution is:  $\frac{P}{Q}$   
(well-approximating rational)



Solution is:  $\frac{p_i}{q_i}$

$$1 \quad \frac{P}{Q} \quad 1 \quad \frac{p_i}{q_i}$$

$$a = \frac{\frac{1}{2Q^2} \frac{P}{q} + \frac{1}{2q^2} \frac{P}{Q}}{\frac{1}{2Q^2} + \frac{1}{2q^2}}$$

$$= \frac{Pq + PQ}{q^2 + Q^2}$$

$$b = \frac{Pq + P_1 q_1}{q^2 + q_1^2}$$

$$c = \frac{P_1 q_1 + PQ}{q_1^2 + Q^2}$$

$$b-a = \frac{q^2 - qQ - Q^2}{(q^2 + Q^2)(q^2 + q_1^2)}$$

Let  $S = \frac{a}{Q} (\geq 1)$ . Then  $S^2 - S - 1$  has same sign as  $b-a$ .

$$\text{Case 1 } a < b, \quad \text{so} \quad S > \frac{\sqrt{5}-1}{2}$$

$$\begin{aligned} \left| \frac{P}{Q} - \alpha \right| &= \frac{P}{Q} - \alpha \leq \frac{P}{Q} - a \\ &= \frac{P}{Q} - \frac{Pq + PQ}{q^2 + Q^2} \\ &= \frac{q}{Q(q^2 + Q^2)} = \frac{S}{S^2 + 1} \cdot \frac{1}{Q^2} \\ &< \frac{1}{\sqrt{5} Q^2} \end{aligned}$$

$$\text{Case 2 } b < a, \quad \text{so} \quad S < \frac{1}{2}(\sqrt{5} + 1)$$

Case 2  $b < a$ , so  $s < \frac{1}{2}(\sqrt{s} + 1)$

$$\begin{aligned} \left| \frac{p_1}{q_1} - \alpha \right| &\leq \frac{p_1}{q_1} - b \quad \text{since } c \text{ is closer to } \frac{p_1}{q_1} \text{ since it's} \\ &= \frac{s(s+1)}{s^2 + (s+1)^2} - \frac{1}{q_1^2} \quad \text{higher on} \\ &< \frac{1}{\sqrt{s} q_1^2} \quad \text{the circle.} \end{aligned}$$

$$SL_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : ad - bc = \pm 1, a, b, c, d \in \mathbb{Z} \right\}.$$


Def  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} z := \frac{az + b}{cz + d}$


$$\text{Then } \left( \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} \right) z = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \left( \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} z \right).$$

Prop  $SL_2(\mathbb{Z})$  acts on Ford circles

In particular,  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} C_{\frac{p}{q}} = C_{\frac{ap+bx}{cq+dy}}$

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in SL_2(\mathbb{Z}), \quad \text{but } \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} C_{\frac{1}{5}} = C_{\frac{1}{5}}$$

so  $C_{\frac{1}{5}} =$  







Prop if we let  $S_{\frac{p}{q}, \frac{p'}{q'}}$  be semicircle joining  $\frac{p}{q}$   $\left| \frac{p'}{q'}\right.$  at right angles,


$S_{\frac{p}{q}, \frac{p'}{q'}}$  goes through  $C_{\frac{p}{q}} \cap C_{\frac{p'}{q'}}$

all of these semicircles are straight lines in hyperbolic geometry.

$SL_2(\mathbb{Z})$  actions preserve these lines.

The Farey Diagram is a "lattice"






### Exercises:

- 1: Find the total area of the Ford Circles  
on the unit interval  $[0,1]$ . (use Ch 17 from Hardy).

Freestyle: Ford spheres w/ Complex Rationals?

2:



Prove this construction  
gives Farey sequences  
(or fix it + prove)

- 3: Use the Cayley transform  $\frac{z+i}{z-i}$

to parametrize Pythagorean triples.

(it takes  $\mathbb{R}$  to boundary of circle)